



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

posteriorly by a quite definite route and rate of extension. The ventral moult is completed before the dorsal. This pelage is somewhat longer and coarser, with a distinct color effect dorsally varying from umber to sepia, and due to increased yellow pigment in the intermediate band of the hair.

The young mice were etherized and the hair plucked out over certain areas, without injuring the skin. Where the juvenal hair was removed it was replaced directly by the post-juvenal. The artificial removal of hair modified the normal sequence of appearance of post-juvenal pelage over the body quite definitely. Usually this modification was confined to the regions actually depilated; but not always. It sometimes influenced the succession at a little distance.

A precocious appearance of the post-juvenal pelage may be induced by removing the juvenile hair.

Restoration takes place in removed adult pelage, and occurs irrespective of seasons. It is restored somewhat more rapidly when hair is plucked out than when it is merely cut. Light appears to have no influence in developing the differences of color in dorsal and ventral surfaces.

EXPERIMENTS ON PROTECTIVE COLORATION

Young (Jour. Exp. Zool. May 1916) reports experiments in which crows, hawks, owls, chickens, prairie chickens, grackles, kingbirds and martins are used as preys, and amphibians, small mammals and insects as prey. The work was done in cages, and varying backgrounds, which contrasted with and concealed the prey were used. He concludes; (1) that protective resemblance is effective in protecting motionless animals from attacks by caged birds; and (2) stillness is probably a more important factor than color in protecting animals from their foes.

COLOR DISCRIMINATION AND ASSOCIATION IN FISHES

White (Jour. Exp. Zool., Feb. 20, 1919) concludes from experiments on mudminnows and sticklebacks that they were able to discriminate, in differing degrees, such colors as red and green, and that the discrimination is based on wave length and not upon intensity. Their power of discrimination is less than that of man. Effective associations between certain colors and behavior were shown by their learning to leap out of the water for food announced by various colors. In a similar way associated actions were shown